Model of Kinetics

The Collision Theory of Chemical Kinetics

The dependence of number of collisions on concentration



doubling the concentration of both reactants

quadrupled the number of collisions

2nd order relationship with respect to the reactants

The dependence of number of collisions on concentration



higher the concentration of reactants

the higher the number of collisions

faster the reaction rate

Collision Theory

in order to react two molecules must encounter one another (collide)



Collision Theory

in order to react two molecules must encounter one another (collide)



Activated complex

A short lived transient species

Collision Theory

in order to react two molecules must encounter one another (collide)



products

Collision Theory

But not all collisions result in reaction



reaction rate is much smaller than the calculated collision frequency ?

Svante Arrhenius

in order to react, the colliding molecules must possess enough kinetic energy to overcome the repulsive and bonding forces of the reactants

activation energy (E_a) the minimum energy required for a chemical reaction to occur

Temperature and Rate of Reaction

At a certain temperature only a certain fraction of the molecules possess enough energy to create effective collisions.



Temperature and Rate of Reaction

effective collisions increase exponentially with an increase in temperature



Temperature and Rate of Reaction

An increase in temperature increases the rate of almost all chemical reactions.

Faster moving particles collide more often and with greater kinetic energy.

Activated complex

the species temporarily formed by the reactant molecules as a result of the collision before they form product



Activated complex

Consider a hypothetical one-step reaction (exothermic)



activated complex

A--------B--------C

Potential energy

activation energy

ΔH

$$\mathrm{A} + \mathrm{B{-}C} \longrightarrow \mathrm{A{-}B} + \mathrm{C}$$

progress of reaction →

Consider a hypothetical one-step reaction (endothermic)



activated complex

A--------B--------C

Potential energy

activation energy

ΔH

$$\mathbf{A} + \mathbf{B{-}C} \longrightarrow \mathbf{A{-}B} + \mathbf{C}$$

progress of reaction →

Molecules must collide with proper geometry



Activated complex



No reaction

Reaction rate is directly proportional to:

- **the number of collisions per second**

  with the proper orientation

- **the fraction of collisions that possess sufficient energy for reaction**