

The Kinetic-Molecular Theory of Gases

kinetic-molecular theory of gases

Originated with Ludwig Boltzmann and James Clerk Maxwell in the 19th century

Explains gas behavior on the basis of the motion of individual particles

Is based on the following assumptions:


(1) A gas is composed of molecules that are separated from each other by distances far greater than their own dimensions.

(2) Gas molecules are in constant motion in random directions, and they frequently collide with one another.

the collisions are completely elastic.


Kinetic-Molecular Theory

Pressure inside container comes from collisions of gas molecules with container walls



This is a model that aids in our understanding of what happens to gas particles as environmental conditions change.





The distances traveled by a single gas molecule between successive collisions. Each change in direction indicates a collision with another molecule.

(3) Gas molecules exert neither attractive nor repulsive forces on one another.

(4) The average kinetic energy of the molecules is proportional to the temperature of the gas in kelvins.

Application to the Gas Laws

- **Compressibility of gases**
- **Boyle's law**
- **Charles's law**
- **Avogadro's law**
- **Dalton's law of partial pressure**

• Compressibility of gases

Recall assumption 1 :

(1) A gas is composed of molecules that are separated from each other by distances far greater than their own dimensions.

Since molecules in the gas phase are separated by large distances, the main result of a decrease in the volume available to them is merely an increase in the rate at which they collide with each other

- **Boyle's law**

Pressure is inversely proportional to volume

gas pressure is the result of collisions between molecules and the walls of the container

a decrease in volume causes an increase in the frequency of collisions between the gas molecule and the container

• Charles's law

recall assumption 4 :

(4) The average kinetic energy of the molecules is proportional to the temperature of the gas in kelvins.

raising the temperature raises the average kinetic energy

**Greater
effect on
pressure**

the frequency of collisions increases

the force of collisions increases

- **Avogadro's law**

recall assumption 1:

(1) A gas is composed of molecules that are separated from each other by distances far greater than their own dimensions.

Equal volumes of gases (at the same temperature and pressure) contain equal numbers of molecules

• Dalton's law of partial pressure

recall assumption 3 :

(3) Gas molecules exert neither attractive nor repulsive forces on one another.

Therefore, the pressure exerted by one type of molecule is unaffected by the presence of another gas.

Therefore, the total pressure is given by the sum of the individual gas pressures.

kinetic-molecular theory

the absolute temperature of a gas is a measure of the average kinetic energy of the molecules

$$(\text{KE})_{\text{ave}} \propto T \qquad (\text{KE})_{\text{ave}} \propto \frac{1}{2} m \overline{u^2}$$

average mean square value of molecular speed

$$\overline{u^2} = \frac{u_1^2 + u_2^2 + \dots + u_N^2}{N_A}$$

kinetic-molecular theory

the absolute temperature of a gas is a measure of the average kinetic energy of the molecules

$$(\text{KE})_{\text{ave}} \propto T \qquad (\text{KE})_{\text{ave}} \propto \frac{1}{2} m \overline{u^2}$$

average mean square value of molecular speed

$$\overline{u^2} = \frac{u_1^2 + u_2^2 + \dots + u_N^2}{N_A}$$

number of molecules in a mole

$$(\text{KE})_{\text{ave}} = \frac{3}{2} RT \qquad (\text{KE})_{\text{ave}} = N_A \left(\frac{1}{2} m \overline{u^2} \right)$$

Root -mean-square speed

one way to estimate how fast molecules are moving at any temperature

$$\sqrt{\overline{u^2}} = u_{rms}$$

kinetic-molecular theory

the absolute temperature of a gas is a measure of the average kinetic energy of the molecules

$$(\text{KE})_{\text{ave}} = \frac{3}{2} RT \qquad (\text{KE})_{\text{ave}} = N_A \left(\frac{1}{2} m \overline{u^2} \right)$$

$$N_A \left(\frac{1}{2} m \overline{u^2} \right) = \frac{3}{2} RT \qquad \overline{u^2} = \frac{3 RT}{N_A m}$$

$$\sqrt{\overline{u^2}} = \sqrt{\frac{3RT}{N_A m}}$$

Root -mean-square speed

$$u_{rms} = \sqrt{\overline{u^2}} = \sqrt{\frac{3RT}{M}}$$

Where M is the molar mass


Root -mean-square speed is directly proportional to the square root kelvin temperature and inversely proportional to the square root of the molar mass (molecular weight)

Distribution of Molecular Speeds


exhibits what is called a “Maxwell-Boltzmann”

- not all molecules move at same speed
- most of the molecules have speeds close to a “most probable speed”
- small number of molecules have very low speeds
- small number have very high speeds
- distribution flattens as temperature increases
- most probable speed increases as temperature increases


Distribution of molecular speeds as a function of temperature.




Distribution of molecular speeds as a function of temperature.



Distribution of molecular speeds as a function of temperature.



Distribution of molecular speeds as a function of temperature.



Graham's Laws of Diffusion and Effusion


Diffusion

the gradual mixing of molecules of one gas with molecules of another by virtue of their kinetic properties


Effusion

the process by which a gas under pressure escapes from one compartment of a container to another by passing through a small opening

Gaseous effusion. Gas molecules move from a high-pressure region (left) to a low-pressure region through a pinhole.




Gaseous effusion. Gas molecules move from a high-pressure region (left) to a low-pressure region through a pinhole.



Light gas ●

heavy gas ●


**Light gases
effuse more
rapidly than
heavy gases.**




Light gas ●

heavy gas ●

Effusion



Both gases effuse through pores in balloon, but lighter helium gas effuses faster than heavier argon gas



© 2012 Pearson Education, Inc.


The difference in the rates of effusion for helium and nitrogen, for example, explains why a helium balloon would deflate faster.



© 2012 Pearson Education, Inc.

Diffusion

Diffusion is the spread of one substance throughout a space or throughout a second substance.



© 2012 Pearson Education, Inc.



Gases

Graham's Law of Effusion

experimental

$$\frac{\text{rate of effusion of gas 1}}{\text{rate of effusion of gas 2}} = \frac{(\text{MW}_2)^{1/2}}{(\text{MW}_1)^{1/2}}$$

**same expression can be derived from
kinetic-molecular theory**

Example

Under identical conditions gaseous CO_2 and CCl_4 are allowed to effuse through a pinhole. If the rate of effusion of the CO_2 is $6.3 \times 10^{-2} \text{ mol s}^{-1}$, what is the rate of effusion of the CCl_4 ?

Graham's Law of Diffusion

$$\frac{\text{rate of effusion of gas 1}}{\text{rate of effusion of gas 2}} = \frac{(\text{MW}_2)^{1/2}}{(\text{MW}_1)^{1/2}}$$

$$\frac{\text{rate of effusion of CCl}_4}{\text{rate of effusion of CO}_2} = \frac{(44\text{g})^{1/2}}{(153.8\text{g})^{1/2}}$$


$$\frac{\text{rate of effusion of CCl}_4}{6.3 \times 10^{-2} \text{ mol/s}} = \frac{(44\text{g})^{1/2}}{(153.8\text{g})^{1/2}}$$

$$\frac{\text{rate of effusion of CCl}_4}{6.3 \times 10^{-2} \text{ mol/s}} = \frac{(44\text{g})^{1/2}}{(153.8\text{g})^{1/2}}$$

$$\text{rate of effusion CCl}_4 = \frac{(44\text{g})^{1/2}}{(153.8\text{g})^{1/2}} \times 6.3 \times 10^{-2} \text{ mol/s}$$

$$= 3.4 \times 10^{-2} \text{ mol/s}$$

Deviation from Ideal Behavior



Plot of PV/RT versus P of 1 mole of a gas at 0°C .

Modifications to Ideal gas law


$$P_1 = \frac{nRT}{V}$$

but volume available to the gas is smaller than V because gas molecules consume some of the space ; this makes the pressure exerted by the gas (P_1) larger than the ideal pressure

$$P_1 = \frac{nRT}{V - nb}$$


(b is a constant characteristic of a particular gas)

Deviations from Ideal Behavior




Low pressure

© 2012 Pearson Education, Inc.




High pressure



Ideal gas

© 2012 Pearson Education, Inc.



Real gas

The assumptions made in the kinetic-molecular model (negligible volume of gas molecules themselves, no attractive forces between gas molecules, etc.) break down at high pressure and/or low temperature.



Modifications to ideal-gas law

but attractive forces between gas molecules make the observed pressure less than that of an ideal gas; subtract a correction factor

$$P_{\text{obs}} = \frac{nRT}{V - nb} - a \frac{n^2}{V^2}$$

(**a** is a constant characteristic of a particular gas)

Van der Waal's equation

van der Waal's constants

	a (atm-L²-mol⁻²)	b L-mol⁻¹
He	0.034	0.0237
Ne	0.211	0.0171
Ar	1.34	0.0322
H₂O	5.46	0.0305

kinetic-molecular theory

because **kinetic energy = $mu^2/2$** , what is important is not the average (mean) velocity but rather the average u^2

$$\text{kinetic energy} = \frac{1}{2} mu^2$$